

## ***Prediction and control of the distraction osteogenesis course. Analytical review***

**A.M. Aranovich, M.V. Stogov, E.A. Kireeva, T.I. Menshchikova**

Russian Ilizarov Scientific Centre for Restorative Traumatology and Orthopaedics, Kurgan, Russian Federation

This review analyzes and assesses the existing methods and approaches to prediction and control of the course of distraction osteogenesis (DO). The analysis of the literature revealed few works that recommended specific predictors or methods for prognosis of the course of distraction osteogenesis at the stages of limb lengthening. The authors identified some diagnostic criteria for assessing the distraction regenerate as potential criteria for predicting its development and maturation. It was found that all available predictors and potential diagnostic criteria for assessing the state of the distraction regenerate in clinical practice are used to further correct the distraction regime (respectively, at the stage of distraction) and to determine the timing of the removal of the apparatus, as well as prognosis of recurrence, fracture, and deformity of the regenerate in the non-apparatus period. It was shown that all known diagnostic methods can be applied for the assessment and prediction of the DO course: radiological, physiological, ultrasound diagnostics, laboratory tests. It is stated that a quantitative assessment of the informative value of most of the known predictors of DO disorders is necessary from the point of view of the evidence-based medicine. Difficulties and problems of the development and application of prognostic tests for assessing DO are described. The directions to the development of this topic are proposed. Predicting the DO course is an essential element for monitoring the tissue repair of the segment under lengthening. Prediction and subsequent prophylaxis of DO disorders is a promising solution for optimizing and improving the quality of treatment of patients with orthopedic diseases by using the Ilizarov method of transosseous distraction osteosynthesis.

**Keywords:** distraction osteosynthesis, Ilizarov method, limb lengthening, prediction

The method of transosseous distraction osteosynthesis (TDO), the fundamental justification and practical use of which were developed by G.A. Ilizarov, enables to achieve significant lengthening of long bones and simultaneous correction of their deformities [1–2]. One of the identified problems in its application is control and management of distraction osteogenesis (DO) [3], which include the diagnosis of the state of distraction regenerate (DR). Mainly radiographic study is used for this purpose followed by, if necessary, correction of treatment tactics. Such a practice, in fact, is just stating the fact and does not allow predicting and preventing possible disturbances in the DO course [4–5]. To solve this problem, an analysis of existing methods for predicting the DO state in clinical practice is necessary. The development of this direction, in our opinion, would enable to qualitatively improve the results of TDO treatment. In addition, this logically fits into the current preventive, predictive and personalized paradigm in contemporary medicine (or 4P medicine) [6].

In connection with the tasks outlined, we conducted an analytical review of the available literature with the aim of assessing the development of the topic on the possibilities and methods for predicting the DO course

in clinical practice with the use of the TDO method.

*Literature search strategy* Open electronic databases of scientific literature PubMed and eLIBRARY were searched for with the keywords: *distraction osteosynthesis, distraction osteogenesis, prognosis, prediction, Ilizarov method, Ilizarov* (in Russian and English versions). The search used separate words and their combination.

To analyze and evaluate the literature data, criteria were determined for including and excluding the sources into the analytical study.

### *Inclusion criteria*

1. Full-text sources or structured abstracts with specific quantitative data
2. Clinical studies indicating that patients were treated using TDO techniques
3. In the sources, distraction regenerate should be evaluated using the characteristics described by quantitative findings

### *Exclusion criteria*

1. Case reports, abstracts of presentations
2. Studies with "duplication" (similar study protocol, groups and number of patients, etc.). Among "duplicate" articles, a more recent source was chosen.

## RESULTS

Literature analysis revealed a small number of works that recommended specific predictors or methods for predicting the course of distraction osteogenesis at the stages of limb bones lengthening with the TDO method.

Therefore, we also included in the analysis those works where the authors proposed quantitative criteria for assessing DO, which are potential predictors for the DO.

A total of 40 sources were included in the analysis [7–46].

The assessment shows that all available predictors and potential diagnostic criteria for evaluating the DO in clinical practice are used mainly for two purposes:

- to correct the distraction mode. In clinical practice, after specific signs indicating a disorder or an increase in the risk of possible DO disorders, as a rule, correction of the distraction regime and/or the use of any additional means (pharmacological correction, physiotherapeutic or other means) should follow. A positive outcome for such preventive measures is achievement of the magnitude of lengthening planned;
- to determine the time-point for removal of the apparatus, the prognosis of recurrence, fractures and regenerate deformity in the period after the removal of the apparatus. Preventive measures at this stage, as

a rule, are associated with procedures for stimulating regeneration maturation, and their positive outcome is a decrease in the frequency of recurrence, deformities, fractures of the regenerate.

Based on this, we grouped the literature published depending on the goals of prognosis.

The methods for DO assessment and prediction in the period of distraction are summarized in Table 1. The table shows that practically all existing diagnostic methods can be applied for the purpose of assessing and predicting the DO course: radiological, ultrasound, laboratory and physiological tests. At the same time, the analysis of these works suggests that laboratory tests (possibility of prediction before surgical treatment and during the first 3–4 days of distraction) and ultrasound diagnosis (the first weeks of distraction) can be classified as the earliest preclinical and pre-roentgenological predictors of DO disturbance.

To assess DR in order to predict abnormalities after removal of the apparatus, the methods of radiological assessment were mainly used: CT, MRI, and bone mineral density (BMD) (Table 2). There are methods of laboratory assessment and prediction of the DR maturation in the period of fixation and in the period after removal of the apparatus.

Table 1

Methods and criteria for assessing and predicting the state of distraction regenerate during the distraction period

Method	Criteria, signs, objects of control	Reference list number
Radiography*	Quantitative DO evaluation, zonography, evaluation of optical density	7, 8, 9, 10
	Height of the connective tissue layer of distraction regenerate	11, 12
	DR shape	13
Ultrasound	DR Echographic signs	14, 15, 16, 17, 18
Physiological	Podography	19
	Thermography	20
Laboratory	Immunogram	21, 22
	Growth factors	23, 24, 25, 26, 27
	Blood biochemistry	28, 29
	Hemostasis	30
	Hematology	31

Note: \* – only quantitative signs were taken into account, qualitative (descriptive) signs were not taken into account

Table 2

Methods and criteria for assessing and predicting the state of distraction regenerate at the fixation stage and in the non-apparatus period

Diagnosis method	Criteria, signs, objects of control	Reference list number
Radiography	Quantitative DO evaluation, zonography, evaluation of optical density	7, 8, 9, 32, 33, 34, 35, 36
	DR Density, diameter, shape	37, 38, 39, 40
	DK BMD	41, 42
MRI	Density of the cortical plate	43
Laboratory test	Blood biochemistry	44, 45, 46

Thus, the data presented allow us to conclude that, there are enough methods for DR assessment to control the DO course. Many of these methods are quite suitable for solving the problems of the DO development prognosis.

However, from the perspective of evidence-based medicine, for most of the tests presented above, a quantitative assessment of the information on these predictors is necessary (calculation

of the odds ratio, sensitivity, specificity). The works in which this quantitative procedure was performed for the proposed criteria and tests are single; all sources are presented in Table 3. It is obvious that the data presented contain tests with a rather high predictive value [38, 40]. However, there are very few validated tests for predicting the DO course during lengthening of limb bones with the Ilizarov method.

Table 3

Value of some criteria for assessing the state of distraction regenerate at the stages of surgical treatment

Method	Predictor	Test value*	Reference
Radiography, DR shape	DR deformity	OR = 5.4 (95 % CI: 2.4-12.4)	[13]
Radiography, DR shape	DR fracture	OR = 19.3 (95 % CI: 2.9-128.0)	[38]
Radiography, DR density and diameter	DR deformity	Sensitivity 93.3 %; Specificity 83.2 %	[40]
Immunogram	DO delay	Prognostic accuracy 81.2 %	[21]

Note: \* - as indicated in the sources; 95 % CI – 95 % confidence interval

### DISCUSSION

The analysis of the literature data shows that currently there are enough methods and means for DR control, but the validated methods and criteria for predicting its state are single.

In addition to absence of reliable methods for predicting DO course, applicable in practice, there are a number of other difficulties in the development of this direction of diagnosis:

a) laboratory tests that seem attractive due to the possibility of early diagnosis are currently not unified and verified, therefore they are used only for research purposes;

b) it is necessary to distinguish between the concepts of diagnosing the current DR state and of its further condition. In practice, it is important for the clinician to understand that the prognosis is based on current diagnostic signs, which only with a certain probability can indicate possible disturbances in the further DO course.

c) the task which follows from the previous paragraph: what are the values of the sensitivity of the tests (odds ratio, specificity, etc.) that are acceptable for predicting adverse events or outcomes when applying the TDO method in clinical practice.

Therefore, in order to improve technologies, methods and prediction criteria for the tasks of DO control, we have identified a number of directions for their development.

1. The search for new sensitive predictors, mainly laboratory ones (molecular genetic and metabolic

studies, etc.).

2. Mandatory unification and validation of potential prognostic tests.

3. Development of comprehensive assessment criteria and DO prediction using several study methods. The application of such approaches was found in a number of works [47, 48].

4. Expansion of prognostic abilities due to the development of technical methods for evaluating, analyzing and prognosis (mathematical modeling, neural networks, software and specialized “diagnostic gadgets” for the Ilizarov apparatus, etc.). Several approaches to the development of this direction were reported in literature sources [49–52].

It is worth noting that the DR evaluation is not the only objective of control and prognosis during surgical lengthening of limb bones with the Ilizarov method. In particular, numerous studies have shown the need and the ability to assess and predict the condition of skeletal muscles and nerves of the segment under lengthening [53–57], blood flow in the segment [58], as well as the prognosis of possible inflammatory reactions associated with the presence of metal implants [59]. Therefore, a comprehensive assessment and prediction of the DO course requires a systematic study considering the state of periosseous organs.

Thus, in terms of developing the direction of DO control and management in the short term, in our opinion, the priority is to validate the known

criteria in order to determine their prognostic value. In the long term (due to the expansion of the number of probable prognostic tests, development of a comprehensive assessment of DR and paraosseous

tissues), the solution of the problems requires creation of an automated (on-line) system for monitoring and predicting the DO course. Moreover, development of such a system is currently quite possible technically.

#### CONCLUSION

Predicting the DO course is a necessary element in solving the problems of monitoring the tissue repair in the segment lengthened with TDO method. Prediction and subsequent prevention of DO disorders in clinical practice are a promising approach for optimizing and

improving the quality of treatment and its outcomes in orthopedic patients who are treated with the Ilizarov method. However, this area of study currently requires significant, primarily technical and statistical development.

#### REFERENCES

- Gubin A., Borzunov D., Malkova T. Ilizarov method for bone lengthening and defect management. Review of contemporary literature. *Bull. Hosp. Jt. Dis.*, 2016, vol. 74, no. 2, pp. 145-154.
- Gubin A.V., Borzunov D.Y., Marchenkova L.O., Malkova T.A., Smirnova I.L. Contribution of G.A. Ilizarov to bone reconstruction: historical achievements and state of the art. *Strategies Trauma Limb Reconstr.*, 2016, vol. 11, no. 3, pp. 145-152. DOI: 10.1007/s11751-016-0261-7.
- Gubin A.V., Borzunov D.Y. Paradigma Ilizarova [Ilizarov's paradigm]. *Genij Ortopedii*, 2012, no. 4, pp. 5-9. (in Russian)
- Borzunov D.Y., Shastov A.L. «Ishemicheskii» distraktsionnyi regenerat pri zameshchenii defektov dlinnykh kostei po G.A. Ilizarovu. Variant resheniia problemy ["Ischemic" distraction regenerated bone when filling long bone defects according to G.A. Ilizarov. Problem solution]. *Vrach-Aspirant*, 2013, vol. 60, no. 5.2, pp. 257-265. (in Russian)
- Urev G.A., Boreiko S.B., Stepuro L.I. Rentgen-ultrazvukovye paralleli v otsenke sostoianiia distraktsionnogo regenerata pri udlinenii konechnosti [Roentgen-ultrasonic parallels in the evaluation of the regenerated bone condition during limb lengthening]. *Meditinskii Zhurnal*, 2008, no. 1 (23), pp. 74-76. (in Russian)
- Paltsev M.A., Belushkina N.N., Chaban E.A. 4P-meditcina kak novaia model zdravookhraneniia v Rossiiskoi Federatsii [4P-medicine as a new model of health service in the Russian Federation]. *ORGZDRAV: novosti, mneniia, obuchenie. Vestnik VShOUZ*, 2015, no. 2, pp. 48-54. (in Russian)
- Diachkova G.V., Mikhailov E.S., Yerofeyev S.A., Nizhechik S.A., Korabelnikov M.A. Kachestvennye i kolichestvennye pokazateli rentgenologicheskoi otsenki distraktsionnogo regenerata [Qualitative and quantitative indices of roentgenological assessment of a distraction regenerate bone]. *Genij Ortopedii*, 2003, no. 4, pp. 11-14. (in Russian)
- Popkov A.V., Aborin S.A., Gorevanov E.A., Klimov O.V. Analiz opticheskoi plotnosti rentgenograficheskogo izobrazheniia distraktsionnogo kostnogo regenerata bedrennoi kosti pri udlinenii vrozhdenno ukorochennogo bedra metodom bilokalnogo distraktsionnogo osteosinteza [The analysis of the optical density of the x-ray image of the femoral distraction regenerate bone in the process of lengthening of congenitally shortened femur using the technique of bifocal distraction osteosynthesis]. *Genij Ortopedii*, 2003, no. 4, pp. 21-23. (in Russian)
- Atanasov N., Poposka A., Samardziski M., Kamnar V. Analysis of radiographic bone parameters throughout the surgical lengthening and deformity correction of extremities. *Pril. (Makedon Akad. Nauk Umet. Odd. Med. Nauki)*, 2014, vol. 35, no. 3, pp. 175-183.
- Catagni M.A., Guerreschi F., Lovisetti L. Distraction osteogenesis for bone repair in the 21st century: lessons learned. *Injury*, 2011, vol. 42, no. 6, pp. 580-586. DOI: 10.1016/j.injury.2011.04.004.
- Dovgalevich I.I. Vnechagovyi kompressionno-distraktsionnyi osteosintez pri lechenii infitsirovannykh defektov dlinnykh trubchatykh kostei [Extrafocal compression-distraktion osteosynthesis in the treatment of infected defects of long tubular bones]. *Voennaia Meditsina*, 2017, no. 1 (42), pp. 20-23. (in Russian)
- Kaliakina V.I. *Sposob distraktsionnogo osteosinteza* [The way of distraction osteosynthesis]. Patent RF, no. 2102022, 1991. (in Russian)
- Li R., Saleh M., Yang L., Coulton L. Radiographic classification of osteogenesis during bone distraction. *J. Orthop. Res.*, 2006, vol. 24, no. 3, pp. 339-347. DOI: 10.1002/jor.20026.
- Eskin N.A., Dorokhin A.I., Pripisnova S.G., Dambinimaev A.V. Vozmozhnosti ultrazvukovogo metoda issledovaniia v otsenke zrelosti distraktsionnogo regenerata pri udlinenii dlinnykh kostei nizhnikh konechnosti [The possibilities of ultrasound research method in the assessment of distraction regenerated bone maturity during lengthening long bones of the lower limbs]. *Biomeditsinskaia Radioelektronika*, 2011, no. 12, pp. 65-72. (in Russian)
- Menshchikova T.I., Aranovich A.M. Osobennosti strukturnogo sostoianiia kostnogo regenerata u bolnykh akhondroplaziei i s vrozhdennoi varusnoi deformatsiei goleni (ultrazvukovoe issledovanie) [Peculiar properties of the structural condition of regenerated bone in patients with achondroplasia and with congenital varus deformity of the leg (An ultrasound study)].



- Vestnik Travmatologii i Ortopedii im. N.N. Priorova*, 2013, no. 1, pp. 68-73. (in Russian)
16. Menshchikova T.I., Aranovich A.M., Dendiberia E.V., Novikov K.I. Ultrasonograficheskie priznaki nizkoi osteogennoi aktivnosti pri udlinenii goleni [Ultrasonographic signs of low osteogenous activity during leg lengthening]. *Genij Ortopedii*, 2004, no. 3, pp. 45-48. (in Russian)
  17. Menshchikova T.I., Borzunov D.Y., Dolganova T.I. Ultrazvukovoe skanirovanie distraktsionnogo regenerata pri polilokalnom udlinenii otlomkov u bolnykh s defektami dlinnykh kostei [Ultrasound scanning of distraction regenerated bone during polylocal fragment lengthening in patients with long bone defects]. *Khirurgiia. Zhurnal im. N.I. Pirogova*, 2014, no. 3, pp. 20-24. (in Russian)
  18. Poposka A., Atanasov N., Dzoleva-Tolevska R. Use of ultrasonography in evaluation of new bone formation in patients treated by the method of Ilizarov. *Prilozi*, 2012, vol. 33, no. 1, pp. 199-208.
  19. Dolganova T.I., Novikov K.I., Kolesnikova E.S. Diagnosticheskaia znachimost podografii pri planirovanii kosmeticheskogo udlineniia goleni [Diagnostic significance of podography when planning cosmetic leg lengthening]. *Fundamentalnye Issledovaniia*, 2014, no. 7-3, pp. 482-486. (in Russian)
  20. Morasiewicz L., Dudek K., Orzechowski W., Kulej M., Stepniewski M. Use of thermography to monitor the bone regenerate during limb lengthening – preliminary communication. *Ortop. Traumatol. Rehabil.*, 2008, vol. 10, no. 3, pp. 279-285.
  21. Belokhvostikova T.S., Mikhalevich I.M. Immunologicheskoe prognozirovanie narushenii regeneratsii kostnoi tkani [Immunological prediction of bone tissue regeneration disorders]. *Sibirskoe Meditsinskoe Obozrenie*, 2018, no. 3 (111), pp. 98-104. (in Russian)
  22. Leonova S.N., Kameka A.L., Rekhov A.V. Primenenie novykh tekhnologii v gnoinoi osteologii [The use of new technologies in purulent osteology]. *Biulleten VSNTs SO RAMN*, 2011, no. 4-1 (80), pp. 110-114. (in Russian)
  23. Vykhovanets E.P., Sakulin N.V., Luneva S.N., Nakoskina N.V., Klimov O.V. Faktory rosta semeistva TGF- $\beta$  v krovi bolnykh akhondroplaziei na etapakh ortopedicheskogo lecheniia [Growth factors of TGF- $\beta$  family in the blood of patients with achondroplasia at the stages of orthopedic treatment]. *Patologicheskaiia Fiziologiia i Eksperimentalnaia Terapiia*, 2018, vol. 62, no. 2, pp. 70-76. (in Russian) DOI: 10.25557/0031-2991.2018.02.70-76.
  24. Compton J., Fragomen A., Rozbruch S.R. Skeletal Repair in Distraction Osteogenesis: Mechanisms and Enhancements. *JBJS Rev.*, 2015, vol. 3, no. 8, 01874474-201508000-00002. DOI: 10.2106/JBJS.RVW.N.00107.
  25. Dhaliwal K., Kunchur R., Farhadieh R. Review of the cellular and biological principles of distraction osteogenesis: An in vivo bioreactor tissue engineering model. *J. Plast. Reconstr. Aesthet. Surg.*, 2016, vol. 69, no. 2, pp. e19-e26. DOI: 10.1016/j.bjps.2015.11.003.
  26. Stogov M.V., Luneva S.N., Novikov K.I. Growth factors in human serum during operative tibial lengthening with the Ilizarov method. *J. Orthop. Res.*, 2013, vol. 31, no. 12, pp. 1966-1970. DOI: 10.1002/jor.22454.
  27. Stogov M.V., Tushina N.V., Emanov A.A. Serum Concentration of Growth Factors in Dogs under Different Conditions of Distraction Osteogenesis. *Bull. Exp. Biol. Med.*, 2015, vol. 160, no. 2, pp. 213-215. DOI: 10.1007/s10517-015-3131-2.
  28. Kovinka M.A., Stogov M.V., Tushina N.V., Gofman F.F. Dinamika biokhimicheskikh pokazatelei syvorotki krovi u patsientov s posttravmaticheskimi ukorocheniami kostei konechnosti v protsesse lecheniia metodom Ilizarova [The dynamics of blood serum biochemical values in patients with posttraumatic limb bone shortenings in the process of treatment by the Ilizarov method]. *Genij Ortopedii*, 2011, no. 4, pp. 35-38. (in Russian)
  29. Talashova I.A., Kononovich N.A., Tushina N.V. Biokhimicheskie markery kosteobrazovaniia i vospaleniia u sobak pri udlinenii kostei goleni po Ilizarovu [Biochemical markers of osteogenesis and inflammation in dogs when lengthening leg bones according to Ilizarov]. *Veterinariia Kubani*, 2016, no. 5, pp. 21-24. (in Russian)
  30. Aranovich A.M., Trofimova E.V., Novikov K.I. *Sposob otsenki techeniia distraktsionnogo osteosinteza* [The way to assess distraction osteosynthesis process]. Patent RF, no. 2291432, 2005. Biul., 2007, no. 1. (in Russian)
  31. Bazarnyi V.V. Laboratornyi monitoring distraktsionnogo kosteobrazovaniia [Laboratory monitoring of distraction osteogenesis]. *Klinicheskaiia Laboratornaia Diagnostika*, 1999, no. 6, pp. 16-17. (in Russian)
  32. Diachkov K.A., Dyachkova G.V., Aleksandrov Iu.M. *Sposob opredeleniia stepeni rezorbtsii kortikalnoi plastinki kosti posle distraktsionnogo udlineniia konechnosti* [The way to determine the resorption degree of cortical bone plate after limb distraction lengthening]. Patent RF, no. 2484772, 2011. Biul., 2013, no. 17. (in Russian)
  33. Dyachkova G.V., Diachkov K.A., Larionova T.A., Aranovich A.M. Kachestvo kosti po dannym multisrezovoi kompiuternoii tomografii (MSKT) u bolnykh s ukorocheniem konechnosti do i posle lecheniia [Bone quality by the data of multislice computed tomography (MSCT) in patients with limb shortening before and after treatment]. *Diagnosticheskaiia i Interventsionnaia Radiologiia*, 2018, vol. 12, no. 1, pp. 11-21. (in Russian)
  34. Popkov A.V., Aborin S.A., Gorevanov E.A., Klimov O.V. Rentgenologicheskaiia dinamika reparativnogo osteogeneza pri bilokalnom distraktsionnom osteosinteze vrozhdennno ukorochennogo bedra [The roentgenological dynamics of reparative osteogenesis for bifocal distraction osteosynthesis of congenitally shortened femur]. *Genij Ortopedii*, 2003, no. 1, pp. 64-67. (in Russian)
  35. Kokoroghiannis C., Charopoulos I., Lyritis G., Raptou P., Karachalios T., Papaioannou N. Correlation of pQCT bone strength index with mechanical testing in distraction osteogenesis. *Bone*, 2009, vol. 45, no. 3, pp. 512-516. DOI: 10.1016/j.bone.2009.05.021.

36. Tesiorowski M., Potaczek T., Jasiewicz B., Kacki W., Łokas K. Regeneration formation index – new method of quantitative evaluation of distraction osteogenesis. *Chir. Narzadów Ruchu Ortop. Pol.*, 2009, vol. 74, no. 3, pp. 121-126.
37. Devmurari K.N., Song H.R., Modi H.N., Venkatesh K.P., Ju K.S., Song S.H. Callus features of regenerate fracture cases in femoral lengthening in achondroplasia. *Skeletal Radiol.*, 2010, vol. 39, no. 9, pp. 897-903. DOI: 10.1007/s00256-009-0742-6.
38. Kitoh H., Mishima K., Matsushita M., Nishida Y., Ishiguro N. Early and late fracture following extensive limb lengthening in patients with achondroplasia and hypochondroplasia. *Bone Joint J.*, 2014, vol. 96-B, no. 9, pp. 1269-1273. DOI: 10.1302/0301-620X.96B9.33840.
39. Shyam A.K., Singh S.U., Modi H.N., Song H.R., Lee S.H., An H. Leg lengthening by distraction osteogenesis using the Ilizarov apparatus: a novel concept of tibia callus subsidence and its influencing factors. *Int. Orthop.*, 2009, vol. 33, no. 6, pp. 1753-1759. DOI: 10.1007/s00264-008-0660-6.
40. Tirawanish P., Eamsobhana P. Prediction of Callus Subsidence in Distraction Osteogenesis Using Callus Formation Scoring System: Preliminary Study. *Orthop. Surg.*, 2018, vol. 10, no. 2, pp. 121-127. DOI: 10.1111/os.12374.
41. Saran N., Hamdy R.C. DEXA as a predictor of fixator removal in distraction osteogenesis. *Clin. Orthop. Relat. Res.*, 2008, vol. 466, no. 12, pp. 2955-2961. DOI: 10.1007/s11999-008-0514-y.
42. Tselentakis G., Owen P.J., Richardson J.B., Kuiper J.H., Haddaway M.J., Dwyer J.S., Evans G.A. Fracture stiffness in callotaxis determined by dual-energy X-ray absorptiometry scanning. *J. Pediatr. Orthop. B*, 2001, vol. 10, no. 3, pp. 248-254.
43. Diachkov K.A., Korabelnikov M.A., Dyachkova G.V., Aranovich A.M., Klimov O.V. MRT-semiotika distraktsionnogo regenerate [MRI-semiotics of distraction regenerated bone]. *Meditinskaja Vizualizatsiia*, 2011, no. 5, pp. 99-103. (in Russian)
44. Lapynin A.I. Biokhimicheskie metody otsenki v otdalennye sroki reabilitatsii bolnykh metodom upravliaemogo chreskostnogo osteosinteza po povodu osteomieliticheskikh polostei kostei nizhnikh konechnostei [Biochemical methods of assessment in the long-term periods of rehabilitation of patients by the method of controlled transosseous osteosynthesis for osteomyelitic cavities of the lower limb bones]. *Izvestiia Samarskogo Nauchnogo Tsentra Rossiiskoi Akademii Nauk*, 2014, vol. 16, no. 5-5, pp. 1808-1811. (in Russian)
45. Popkov D.A., Kuznetsova L.S., Luneva S.N., Stogov M.V. Vozmozhnosti nagruzochnoi proby laktatom kaltsiia dlia otsenki sostoianiia kaltsii-reguliruiushchei gormonalnoi sistemy pri udlinenii konechnostei [The potentials of the tolerance test with calcium lactate to assess the calcium-regulating hormonal system state during limb lengthening]. *Genij Ortopedii*, 2005, no. 4, pp. 65-68. (in Russian)
46. Tushina N.V., Stogov M.V., Kononovich N.A., Emanov A.A. Prognoz dlitelnosti sozrevaniia distraktsionnogo regenerata [The prediction of the duration of the distraction regenerated bone maturation]. *Travmatologiya i Ortopediia Rossii*, 2012, no. 1 (63), pp. 49-54. (in Russian)
47. Menshchikova T.I., Novikov K.I. Ultrasonograficheskaja i rentgenologicheskaja otsenki strukturnogo sostoianiia distraktsionnogo regenerata bolshebertsovoi kosti [Ultrasound and X-ray assessment of the structural condition of the tibial distraction regenerated bone]. *Travmatologiya i Ortopediia Rossii*, 2005, no. 4 (38), pp. 57-59. (in Russian)
48. Saifutdinov M.S., Diachkov K.A., Korabelnikov M.A. Sopostavitelnyi analiz aktivatsionnykh kharakteristik myshts nizhnikh konechnostei do i posle ikh udlineniiia metodom distraktsionnogo osteosinteza u bolnykh akhondroplaziei s dannymi kompiuternoi tomografii [The comparative analysis of the activation characteristics of lower limb muscles before and after their lengthening by distraction osteosynthesis method in patients with achondroplasia who have the data of computer tomography]. *Genij Ortopedii*, 2006, no. 3, pp. 73-77. (in Russian)
49. Aranovich A.M., Dyachkova G.V., Klimov O.V., Diachkov K.A., Neretin A.C. Metodiki tsifrovogo analiza rentgenologicheskogo izobrazheniia distraktsionnogo regenerata pri udlinenii golenei u bolnykh akhondroplaziei [The techniques of digital analysis of the X-ray image of distraction regenerated bone during leg lengthening in patients with achondroplasia]. *Fundamentalnye Issledovaniia*, 2015, no. 1-6, pp. 1115-1119. (in Russian)
50. Gaidyshev I.P. Programmnoe upravlenie apparatom Ilizarova [Program control of the Ilizarov fixator]. *Genij Ortopedii*, 2005, no. 3, pp. 97-101. (in Russian)
51. Krupchatnikov R.A., Mishustin V.N., Artemenko M.V., Dobrovolskii I.I. Metody diagnostiki i faktory riska prognoza tromboembolii v sistemakh podderzhki priniatiia reshenii [Diagnostic methods and risk factors of thromboembolism prediction in the systems of decision making support]. *Izvestiia Iugo-Zapadnogo Gosudarstvennogo Universiteta. Seriya: Upravlenie, vychislitelnaia tekhnika, informatika. Meditsinskoe Priborostroenie*, 2017, vol. 7, no. 1 (22), pp. 69-81 (in Russian)
52. Simpson A.L., Ma B., Slagel B., Borschneck D.P., Ellis R.E. Computer-assisted distraction osteogenesis by Ilizarov's method. *Int. J. Med. Robot.*, 2008, vol. 4, no. 4, pp. 310-320. DOI: 10.1002/rcs.211.
53. Vinokurova T.S., Garkavenko Iu.E., Brytov A.V. Kliniko-neirofiziologicheskii analiz funktsionalnogo sostoianiia perifericheskoi nervnoi sistemy u detei s posledstviiami gematogenno osteomielita pri udlinenii nizhnei konechnosti metodom distraktsionnogo osteosinteza [Clinical and neurophysiological analysis of the peripheral nervous system functional condition in children with hematogenous osteomyelitis consequences when lengthening the lower limb by the method of distraction osteosynthesis]. *Travmatologiya i Ortopediia Rossii*, 2008, no. 3 (49), pp. 30-34. (in Russian)

54. Grebenyuk L.A., Saifutdinov M.S., Grebeniuk E.B. Vozmozhnosti kompleksnogo ispolzovaniia ekhografii i elektromiografii v otsenke sostoianiia nervno-myshechnogo apparata pri operativnom udlinenii konechnosti [Possibilities of complex using echography and electromyography in the assessment of the neuromuscular system during surgical limb lengthening]. *Sovremennye Problemy Nauki i Obrazovaniia*, 2016, no. 1, pp. 33. (in Russian)
55. Menshchikova T.I., Aranovich A.M. Otsenka rezervnykh vozmozhnostei perednei gruppy myshts goleni u bolnykh akhondroplaziei v protsesse udlineniia (ultrazvukovoe issledovanie) [Evaluation of the reserve capacity of the leg anterior muscle group in patients with achondroplasia during lengthening (An ultrasound study)]. *Vestnik Travmatologii i Ortopedii im. N.N. Priorova*, 2015, no. 3, pp. 13-19. (in Russian)
56. Nogueira M.P., Paley D., Bhave A., Herbert A., Nocente C., Herzenberg J.E. Nerve lesions associated with limb-lengthening. *J. Bone Joint Surg. Am.*, 2003, vol. 85, no. 8, pp. 1502-1510. DOI: 10.2106/00004623-200308000-00011.
57. Novikov K.I., Klimov O.V., Subramanyam K.N., Neretin A.S., Novikova O.S. Myotopography in Tibial Lengthening. *Arch. Bone Jt. Surg.*, 2018, vol. 6, no. 4, pp. 331-334.
58. Shchurov V.A., Popkov A.V., Aranovich A.M. Poliarograficheskii kontrol sostoianiia regionarnogo krovoobrashcheniia pri operativnom udlinenii goleni [Polarographic monitoring of regional circulation state during surgical leg lengthening]. *Genij Ortopedii*, 2015, no. 4, pp. 76-79. (in Russian) DOI: 10.18019/1028-4427-2015-4-76-79.
59. Kutepov S.M., Shlykov I.L., Berdiugina O.V., Berdiugin K.A., Kuznetsova N.L. Prognozirovanie vospalitelnykh oslozhnenii pri vneochagovom osteosinteze [Prediction of inflammatory complications for extrafocal osteosynthesis]. *Vestnik Travmatologii i Ortopedii im. V.D. Chaklina*, 2009, vol. 1, no. 1, pp. 44-46. (in Russian)

Received: 05.03.2019

**Information about the authors:**

1. Anna M. Aranovich, M.D., Ph.D., Professor,  
Russian Ilizarov Scientific Centre for Restorative Traumatology and Orthopaedics, Kurgan, Russian Federation
2. Maksim V. Stogov, Ph.D. of Biological Sciences,  
Russian Ilizarov Scientific Centre for Restorative Traumatology and Orthopaedics, Kurgan, Russian Federation,  
Email: stogo\_off@list.ru
3. Elena A. Kireeva, Ph.D. of Biological Sciences,  
Russian Ilizarov Scientific Centre for Restorative Traumatology and Orthopaedics, Kurgan, Russian Federation
4. Tatiana I. Menshchikova, Ph.D. of Biological Sciences,  
Russian Ilizarov Scientific Centre for Restorative Traumatology and Orthopaedics, Kurgan, Russian Federation,  
Email: tat-mench@mail.ru